

## Recombinant Human ILDR2 ECD-Human Fc

**Catalog Number/Size:** P0841-100 100 μg

P0841-B Bulk

Source: Human ILDR2 (Accession # XP\_016856747.1) extracellular domain (Leu36-Glu201) fused with human IgG1 Fc

produced from HEK293 cells.

Human ILDR2 (Leu36-Glu201)
Accession#XP\_016856747.1

GSGGGG Human IgG1 (Asp104-Lys330)

N-terminal

C-terminal

N-terminal C-te

**Structure:** Disulfide-linked homodimer

**Predicted N-terminal:** Leu 36

**Predicted Molecular** 

Weight:

89.3 kDa (dimer)

Apparent Molecular Weight on SDS-PAGE:

arent Molecular 50.0 kDa, reducing conditions


**Formulation:** 0.22 µm filtered protein solution in PBS

**Storage:** < -20°C

**Estimated Purity:** >95% as determined by SDS-PAGE

Protein Endotoxin Level: Not measured
Protein Aggregation: Not measured

## DATA



Disclaimer: For research use only. Not for use in humans.



## Recombinant Human ILDR2 ECD-Human Fc

**Application:** Biochemical analysis

**Product Description:** ILDR2 (immunoglobulin-like domain-containing receptor 2) has four major isoforms. It is a single-pass type I

transmembrane protein containing an amino terminal immunoglobulin-like domain (IgV like) and a long, carboxy tail (1). It was found to localize in the endoplasmic reticulum and is suggested to be involved in lipid homeostasis and ER stress pathways (2). It has also been associated with type 2 diabetes (1). It belongs to immunoglobulin

superfamily and LISCH7 family (3).

Other Names: Clorf32, "Lisch-like"

References: 1. Dokmanovic-Chouinard, M. et al. (2008) PLoS Genet. 4(7).

2. Watanabe, K. et al. (2013) PLoS ONE. 8(6).

3. Tomohihto, H. et al. (2013) J. Cell. Sci. 126:966.

Disclaimer: For research use only. Not for use in humans.