


SPECIFICATIONS

Catalog Number	C3054
Cell Line Name	Mouse B7-H4-CHO-K1 stable cell line
Accession Number	NP_848709.2
Host Cell	Adherent CHO-K1
Quantity	Two vials of frozen cells (2x10 ⁶ per vial)
Culture Medium	DMEM with 10% FBS, 4µg/ml puromycin
Freezing Medium	90% FBS and 10% DMSO
Storage	Liquid nitrogen

DATA

Detection of mouse B7-H4 expression on mouse B7-H4-CHO-K1 stable cells using an anti-human B7-H4 antibody, followed by staining with PE-anti-mouse IgG antibody.


BACKGROUND

B7-H4, also known as VTCN1 (V-set domain-containing T-cell activation inhibitor 1, B7X, B7H4, and B7S1), belongs to the B7 family of immune regulatory proteins expressed on the surface of antigen-presenting cells (APCs), such as dendritic cells, macrophages, and B cells. Its expression is induced by pro-inflammatory cytokines and can be upregulated in response to immune activation. The main function of B7-H4 is to negatively regulate T-cell-mediated immune responses by inhibiting the proliferation, cytokine secretion, and cell cycle of T cells. B7-H4 expression has been observed in various tumor types, including breast, ovarian, lung, gastric, and pancreatic cancers. Its upregulation in these tumors has been associated with poor prognosis and reduced patient survival. The overexpression of B7-H4 in cancer cells is thought to contribute to immune evasion by suppressing the anti-tumor immune response.

References

- Wang, Jia-Yu, and Wei-Peng Wang. B7-H4, a promising target for immunotherapy." *Cellular immunology* .**347**:104008. 2020.
- Zang, Xingxing et al. "B7x: a widely expressed B7 family member that inhibits T cell activation." *Proceedings of the National Academy of Sciences of the United States of America*. **100**,18: 10388-92. 2003.
- Tringler, Barbara et al. B7-h4 is highly expressed in ductal and lobular breast cancer. *Clinical Cancer Research*. **5**: 1842-8. 2005.

Disclaimer: For research use only. Not for use in humans.